

Tennessee River Basin

Short Creek Embayment Guntersville Reservoir Intensive Basin Survey 2009

GUNM-8: Short Creek immediately upstream of AL Hwy 227 bridge (Marshall Co 34.3645/-86.2169)

BACKGROUND

The Alabama Department of Environmental Management (ADEM) began monitoring lake water quality statewide in 1985, followed by a second statewide survey in 1989. In 1990, the Reservoir Water Quality Monitoring Program (now known as the Rivers and Reservoirs Monitoring Program (RRMP) was initiated by ADEM.

The current objectives of this program are to provide data that can be used to assess current water quality conditions, identify trends in water quality conditions and to develop Total Maximum Daily Loads (TMDLs) and water quality criteria. Descriptions of all RRMP monitoring activities are available in ADEM's 2012 Monitoring Strategy (ADEM 2012).

In 2009, ADEM monitored the Short Creek tributary embayment of Guntersville Reservoir as part of the intensive basin assessment of the Tennessee River under the RRMP. This site was selected using historical data and previous assessments. The purpose of this report is to summarize data collected in the Short Ck embayment (GUNM-8) during the 2009 growing season (Apr-Oct). This is the second intensive basin assessment of the Tennessee River since ADEM began sampling on a basin rotation. Monthly and/or mean concentrations of nutrients [total nitrogen (TN); total phosphorus (TP)], algal biomass/productivity [chlorophyll *a* (chl *a*); algal growth potential testing (AGPT)], sediment [total suspended solids (TSS)], and trophic state [Carlson's trophic state index (TSI)] from 2009 were compared to ADEM's 2003 data and established criteria.

WATERSHED CHARACTERISTICS

Watershed land uses are summarized in Table 1. Short Creek is classified as a *Public Water Supply/Swimming/Fish & Wildlife (PWS/S/F&W)* stream located in the Southern Table Plateaus ecoregion (68d). Based on the 2006 National Land Cover Dataset, land use within the 224 mi² watershed is predominantly agriculture [hay/pasture (45%) and cultivated crops (9%)](Fig. 3). As of October 1, 2013, ADEM has issued a total of 52 NPDES permits within the watershed. Thirteen of those permits are located within 10 mi of the station (Fig. 2).

SITE DESCRIPTION

The Short Ck embayment at GUNM-8 is located northeast of Guntersville, AL. This watershed handles all of the runoff and discharges from the city of Albertville. It is a fairly large embayment flowing into the Tennessee River near Guntersville State Park at river mile 361. Short Ck has a median bottom depth of 10.2 m (Table 2) at the sampling location.



Figure 1. Photo of Short Ck at GUNM-8




Figure 2. Map of Short Ck Embayment of Guntersville Reservoir. Though additional discharges may occur in the watershed (Table 1), only permitted discharges within 10 miles upstream of the station are displayed on the map.

METHODS

Water quality assessments were conducted at monthly intervals, April-October. All samples were collected, preserved, stored, and transported according to procedures in the ADEM Field Operations Division Standard Operating Procedures (ADEM 2009), Surface Water Quality Assurance Project Plan (ADEM 2008a), and Quality Management Plan (ADEM 2008b).

Mean growing season TN, TP, chl *a*, and TSS were calculated to evaluate water quality conditions. Monthly concentrations of these parameters were graphed with discharge data, if available, and ADEM's previously collected data to help interpret the 2009 results. Carlson's TSI was calculated from the corrected chl *a* concentrations.

Table 1: Summary of Watershed GUNM-8

Table 1: Summary of Watershed	GUNM-8		
Basin	Tennessee R		
Drainage Area (mi²)	224		
Ecoregion ^a	68d		
% Landuse			
Open Water	1%		
Developed Open Space	7%		
Low Intensity	<1%		
Medium Intensity	<1%		
High Intensity	<1%		
Barren Land	<1%		
Forest Deciduous Forest	14%		
Evergreen Forest	5%		
Mixed Forest	10%		
Shrub/Scrub	3%		
Herbaceous	1%		
Hay/Pasture	45%		
Cultivated Crops	9%		
Wetlands Woody	0%		
Emergent Herb.	<1%		
#NPDES Permits ^b TOTAL	52		
Construction Stormwater	16		
Industrial General	26		
Industrial Individual	1		
Municipal Individual	4		
Underground Injection Control	5		

a. Southern Table Plateaus




Figure 3. Landuse within the Short Creek watershed at GUNM-8.

RESULTS

The following discussion of results is limited to those parameters which directly affect trophic status or parameters which have established criteria. Results of all water chemistry analyses are presented in Table 2. The axis of the graphs in Fig. 4 were set to maximum values reservoir wide so all embayment reports on the same reservoir could be compared.

The mean growing season TN value was higher in 2009 than in 2003 (Fig. 4). Monthly TN concentrations peaked in April.

The mean growing season TP concentration was slightly lower in 2009 (Fig. 4). Monthly TP concentrations decreased April-August and peaked in September.

In 2009, the growing season mean chl a value was lower than in 2003 (Fig. 4). Monthly chl a concentrations were highest in June and July.

Mean TSI remained eutrophic in 2009 (Fig.4). Monthly TSI in Short Ck was mesotrophic in April, and eutrophic May-October.

The mean growing season TSS value was lower in 2009 than 2003 (Fig. 5). Monthly TSS concentrations were highest in April with a second peak in July after a sharp drop in May.


AGPT results show that GUNM-8 was nitrogen limited in both 2003 and 2009 (Table 3). The mean maximum standing crop (MSC) value was 8.19 mg/L, which is above the 5.0 m/L value that Raschke and Schultz (1987) defined as protective of reservoir and lake systems. The previous MSC value for Short Ck was also above 5 mg/L.

The DO concentration in the GUNM-8 station was below the ADEM criteria limit of 5.0 mg/l at 5.0 ft (1.5 m) in August (ADEM Admin. Code R. 335-6-10-.09) (Fig. 6). All other DO measurements were above the criteria limit.

b. #NP DES permits do wnlo aded from ADEM's NP DES Management System database, Oct 1, 2013.



Figure 4. Mean growing season (2003-2009) and monthly (April-October, 2009) TN, TP, chl a and TSI measured in the Short Creek embayment of Guntersville Reservoir. Vertical axis ranges are set to maximum values reservoir-wide for comparability between embayment reports within the same reservoir.






Figure 5. Mean growing season and monthly TSS measured in the Short Creek embayment of Guntersville Reservoir.

Table 2. Summary of water quality data collected April-October, 2009. Minimum (Min) and maximum (Max) values calculated using minimum detection limits. Median (Med), average (Avg), and standard deviations (SD) values were calculated by multiplying the MDL by 0.5 when results were less than this value.

GUNM-8	N	Min	Max	Med	Avg	SD
Physical						
Turbidity (NTU)	7	3.1	13.0	5.2	6.2	3.2
Total Dissolved Solids (mg/L) ^J	7	66.0	128.0	118.0	104.3	24.6
Total Suspended Solids (mg/L)	7	< 1.0	9.0	5.0	4.6	3.1
Hardness (mg/L)	3	49.2	75.6	70.5	65.1	14.0
Alkalinity (mg/L)	7	17.5	76.1	64.8	56.2	20.2
Photic Zone (m)	7	2.67	4.05	3.01	3.18	0.52
Secchi (m)	7	0.65	1.44	1.15	1.10	0.27
Bottom Depth (m)	8	10.00	10.50	10.20	10.20	0.15
Chemical						
Ammonia Nitrogen (mg/L)	7	< 0.006	0.053	0.007	0.012	0.018
Nitrate+Nitrite Nitrogen (mg/L) ^J	7	0.018	1.395	0.302	0.418	0.490
Total Kjeldahl Nitrogen (mg/L)	7	< 0.089	0.779	0.435	0.417	0.245
Total Nitrogen (mg/L) ^J	7	< 0.346	1.985	0.818	0.835	0.558
Dissolved Reactive Phosphorus (mg/L) ^J	7	0.005	0.066	0.020	0.025	0.021
Total Phosphorus (mg/L)	7	0.047	0.099	0.064	0.067	0.018
CBOD-5 (mg/L)	7	< 2.0	2.0	1.0	1.0	0.0
Chlorides (mg/L)	7	5.4	10.2	6.0	7.0	1.9
Biological						
Chlorophy II a (ug/L)	7	5.34	25.10	15.31	15.44	7.95
Fecal Coliform (col/100 mL) ^J	3	< 1	7	6	5	3

J= one or more of the values is an estimate; N= # samples.

Table 3. Algal growth potential test results (expressed as mean MSC) dry weights of *Selenastrum capricornutum* in mg/L) and limiting nutrient status. MSC values below 5 mg/L are considered to be protective in reservoirs and lakes (Raschke and Schultz 1987).

Year	Mean MSC	Limiting Nutrient
8/20/2003	9.32	NITROGEN
8/18/2009	8.19	NITROGEN

FOR MORE INFORMATION, CONTACT: Gina Curvin, ADEM Environmental Indicators Section 1350 Coliseum Boulevard, Montgomery, AL 36110 (334) 260-2783, gcurvin@adem.state.al.us




Figure 6. Monthly DO concentrations at 1.5 m (5 ft) for Short Ck embayment station of Guntersville Reservoir collected April-October 2009. ADEM Water Quality Criteria pertaining to reservoir waters require a DO concentration of 5.0 mg/L at this depth.

REFERENCES

ADEM. 2008a. Quality Assurance Project Plan (QAPP) for Surface Water Quality Monitoring in Alabama. Alabama Department of Environmental Management (ADEM), Montgomery, AL. 78 pp.

ADEM. 2008b. Quality Management Plan (QMP) for the Alabama Department of Environmental, Alabama Department of Environmental Management (ADEM), Montgomery, AL. 58 pp.

ADEM. 2009. Standard Operating Procedures Series #2000, Alabama Department of Environmental Management (ADEM), Montgomery, AL.

ADEM. 2012. State of Alabama Water Quality Monitoring Strategy June 19, 2012. Alabama Department of Environmental Management (ADEM), Montgomery, AL. 88 pp.http://www.adem.alabama.gov/programs/water/wqsurvey/2012WQMonitoringStrategy

Alabama Department of Environmental Management Water Division (ADEM Admin. Code R. 335-6-10-.09). 2010. Specific Water Quality Criteria. Water Quality Program. Chapter 10. Volume 1. Division 335-6.

Alabama Department of Environmental Management Water Division (ADEM Admin. Code R. 335-6-10-.11). 2010. Water Quality Criteria Applicable to Specific Lakes. Water Quality Program. Chapter 10. Volume 1. Division 335-6.

Carlson, R.E. 1977. A trophic state index. Limnology and Oceanography. 22(2):361-369.

Raschke, R.L. and D.A. Schultz. 1987. The use of the algal growth potential test for data assessment. Journal of Water Pollution Control Federation 59(4):222-227.