

Round Island Creek Embayment Wheeler Reservoir Intensive Basin Survey 2009

WHEL-8: Round Island Creek approx 1.5 mi upstream of confluence with TN R (Limestone Co 34.6986/-87.0507)

BACKGROUND

The Alabama Department of Environmental Management (ADEM) began monitoring lake water quality statewide in 1985, followed by a second statewide survey in 1989. In 1990, the Reservoir Water Quality Monitoring Program (now known as the Rivers and Reservoirs Monitoring Program (RRMP) was initiated by ADEM.

The current objectives of this program are to provide data that can be used to assess current water quality conditions, identify trends in water quality conditions and to develop Total Maximum Daily Loads (TMDLs) and water quality criteria. Descriptions of all RRMP monitoring activities are available in ADEM's 2012 Monitoring Strategy (ADEM 2012).

In 2009, ADEM monitored the Round Island Creek tributary embayment of Wheeler Reservoir as part of the intensive basin assessment of the Tennessee River under the RRMP. This site was selected using historical data and previous assessments. The purpose of this report is to summarize data collected in the Round Island Creek embayment (WHEL-8) during the 2009 growing season (Apr-Oct). This is the second intensive basin assessment of the Tennessee River since ADEM began sampling on a basin rotation. Monthly and/or mean concentrations of nutrients [total nitrogen (TN); total phosphorus (TP)], algal biomass/productivity [chl *a*; algal growth potential testing (AGPT)], sediment [total suspended solids (TSS)], and trophic state [Carlson's trophic state index (TSI)] from 2009 were compared to ADEM's 2003 data and established criteria.

WATERSHED CHARACTERISTICS

Watershed land uses are summarized in Table 1. Round Island Creek is classified as a *Fish & Wildlife (F&W)* stream located in the Eastern Highland Rim ecoregion (71g). Based on the 2006 National Land Cover Dataset, land use within the 52 mi² watershed is predominantly agriculture [hay/pasture (42%) and crops (21%)] (Fig. 3). As of October 1, 2013, ADEM has issued a total of 8 NPDES permits within the watershed. Two of those permits are located within 10 mi of the station (Fig. 2).

SITE DESCRIPTION

The Round Island Creek embayment at WHEL-8 is located just south of Athens, AL. It is a fairly wide and shallow embayment which flows into the Tennessee River near river mile 297. Round Island Creek has a mean bottom depth of 1.65 m (Table 2) at the sampling location.



Figure 1. Photo of Round Island Creek at WHEL-8




Figure 2. Map of Round Island Creek embayment of Wheeler Reservoir. Though additional permits may occur in the watershed (Table 1), only permitted discharges within 10 miles upstream of the station are displayed on the map.

METHODS

Water quality assessments were conducted at monthly intervals, April-October. All samples were collected, preserved, stored, and transported according to procedures in the ADEM Field Operations Division Standard Operating Procedures (ADEM 2009), Surface Water Quality Assurance Project Plan (ADEM 2008a), and Quality Management Plan (ADEM 2008b).

Mean growing season TN, TP, chl *a*, and TSS were calculated to evaluate water quality conditions. Monthly concentrations of these parameters were graphed with ADEM's previously collected data to help interpret the 2009 results. Carlson's TSI was calculated from the corrected chl *a* concentrations.

RESULTS

The following discussion of results is limited to those parameters which directly affect trophic status or parameters which have established criteria. Results of all water chemistry analyses are presented in Table 2. The axis ranges of the graphs in Figs. 4-6 were set to maximum values reservoir wide so all embayment reports on the same reservoir could be compared.

Table 1: Summary of Watershed WHEL-8

Basin		Tennessee R
Drainage Area (mi ²)		52
Ecoregion ^a		71g
% Land use		
Open Water		<1%
Developed	Open Space	5%
	Low Intensity	1%
	Medium Intensity	<1%
Barren Land		<1%
Forest	Deciduous Forest	13%
	Evergreen Forest	2%
	Mixed Forest	1%
Shrub/Scrub		5%
Herbaceous		2%
Hay/Pasture		42%
Cultivated Crops		21%
Wetlands	Woody	7%
	TOTAL	8
# NPDES Permits ^b		
401 Water Quality Certification		1
Construction Stormwater		4
Industrial General		3

a. Eastern Highland Rim

b. #NPDES permits downloaded from ADEM's NPDES Management System database, Oct 1, 2013.




Figure 3. Land use within the Round Island Creek watershed at WHEL-8.

The mean growing season TN value was much higher in 2009 than in 2003 (Fig. 4). Monthly TN concentrations were highest in June and October.

The mean growing season TP concentration was slightly higher in 2009 (Fig. 4). The highest monthly TP concentration was measured in June.

In 2009, the growing season mean chl *a* value was higher than 2003 (Fig. 4). Monthly chl *a* concentrations peaked in August and declined sharply through October.

Mean TSI was eutrophic in 2009 and 2003. Monthly TSI in Round Island Creek was eutrophic April-September (Fig. 4).

The mean growing season TSS value was lower in 2009 than 2003 (Fig. 5). Monthly TSS concentration generally increased April-August and declined through October.

AGPT results show that WHEL-8 was nitrogen limited in both 2003 and 2009 (Table 3). The mean maximum standing crop (MSC) value from both years were above the 5.0 mg/L value that Raschke and Schultz (1987) defined as protective of reservoir and lake systems.

The DO concentration in the WHEL-8 station was above the ADEM criteria limit of 5.0 mg/l at 5.0 ft (1.5 m) in all months (ADEM Admin. Code R. 335-6-10-.09) (Fig. 6).











Figure 4. Mean growing season (2003-2009) and monthly (April-October, 2009) TN, TP, chl a and TSI measured in the Round Island Creek embayment of Wheeler Reservoir. Vertical axis ranges are set to maximum values reservoir-wide for comparability between embayment reports within the same reservoir.





Figure 5. Mean growing season and monthly TSS measured in the Round Island Creek embayment of Wheeler Reservoir.

Table 2. Summary of water quality data collected April-October, 2009. Minimum (Min) and maximum (Max) values calculated using minimum detection limits. Median (Med), mean, and standard deviations (SD) values were calculated by multiplying the MDL by 0.5 when results were less than this value.

WHEL-8	N	Min	Max	Med	Mean	SD
Physical						
Turbidity (NTU)	7	5.0	49.6	23.1	25.0	13.1
Total Dissolved Solids (mg/L) ^J	7	28.0	132.0	76.0	76.9	39.8
Total Suspended Solids (mg/L) ^J	7	< 1.0	33.0	17.0	18.8	11.9
Hardness (mg/L)	3	35.1	60.6	50.4	48.7	12.8
Alkalinity (mg/L)	7	34.0	65.0	54.6	48.9	11.9
Photic Zone (m)	6	1.16	2.36	1.44	1.55	0.43
Secchi (m)	6	0.27	1.36	0.42	0.56	0.40
Bottom Depth (m)	7	1.00	2.36	1.47	1.65	0.48
Chemical						
Ammonia Nitrogen (mg/L)	7	< 0.006	0.014	0.007	0.005	0.002
Nitrate+Nitrite Nitrogen (mg/L)	7	0.036	1.455	0.507	0.520	0.478
Total Kjeldahl Nitrogen (mg/L)	7	0.232	1.194	0.520	0.624	0.344
Total Nitrogen (mg/L)	7	0.329	1.915	1.114	1.143	0.572
Dissolved Reactive Phosphorus (mg/L) ^J	7	0.005	0.069	0.010	0.026	0.027
Total Phosphorus (mg/L)	7	0.029	0.179	0.074	0.091	0.049
CBOD-5 (mg/L)	7	< 2.0	2.0	1.0	1.0	0.0
Chlorides (mg/L)	7	3.3	7.4	5.5	5.4	1.2
Biological						
Chlorophyll a (ug/L)	7	< 0.10	28.48	21.36	16.32	9.83
Fecal Coliform (col/100 mL) ^J	3	5	400	93	166	207

J= one or more of the values is an estimate; N= # samples.

Table 3. Algal growth potential test results (expressed as mean MSC) dry weights of *Selenastrum capricornutum* in mg/L) and limiting nutrient status. MSC values below 5 mg/L are considered to be protective in reservoirs and lakes (Raschke and Schultz 1987).

Year	Mean MSC	Limiting Nutrient
8/20/2003	5.2	NITROGEN
8/18/2009	7.95	NITROGEN

FOR MORE INFORMATION, CONTACT:

Gina Curvin, ADEM Environmental Indicators Section
1350 Coliseum Boulevard, Montgomery, AL 36110
(334) 260-2783, gcurvin@adem.state.al.us




Figure 6. Monthly DO concentrations at mid-depth for Round Island Creek embayment station of Wheeler Reservoir collected April-October 2009. ADEM Water Quality Criteria pertaining to reservoir waters require a DO concentration of 5.0 mg/L at this depth.

REFERENCES

- ADEM. 2008a. Quality Assurance Project Plan (QAPP) for Surface Water Quality Monitoring in Alabama. Alabama Department of Environmental Management (ADEM), Montgomery, AL. 78 pp.
- ADEM. 2008b. Quality Management Plan (QMP) for the Alabama Department of Environmental, Alabama Department of Environmental Management (ADEM), Montgomery, AL. 58 pp.
- ADEM. 2009. Standard Operating Procedures Series #2000, Alabama Department of Environmental Management (ADEM), Montgomery, AL.
- ADEM. 2012. State of Alabama Water Quality Monitoring Strategy June 19, 2012. Alabama Department of Environmental Management (ADEM), Montgomery, AL. 88 pp. <http://www.adem.alabama.gov/programs/water/wqsurvey/2012WQMonitoringStrategy>
- Alabama Department of Environmental Management Water Division (ADEM Admin. Code R. 335-6-10-.09). 2010. Specific Water Quality Criteria. Water Quality Program. Chapter 10. Volume 1. Division 335-6.
- Alabama Department of Environmental Management Water Division (ADEM Admin. Code R. 335-6-10-.11). 2010. Water Quality Criteria Applicable to Specific Lakes. Water Quality Program. Chapter 10. Volume 1. Division 335-6.
- Carlson, R.E. 1977. A trophic state index. *Limnology and Oceanography*. 22(2):361-369.
- Raschke, R.L. and D.A. Schultz. 1987. The use of the algal growth potential test for data assessment. *Journal of Water Pollution Control Federation* 59(4):222-227.